
Frequency of mononuclear diploid cardiomyocytes underlies natural variation in heart regeneration.

Journal: Nat Genet

Publication Year: 2017

Authors: Michaela Patterson, Lindsey Barske, Ben Van Handel, Christoph D Rau, Peiheng Gan, Avneesh Sharma, Shan Parikh, Matt Denholtz, Ying Huang, Yukiko Yamaguchi, Hua Shen, Hooman Allayee, J Gage Crump, Thomas I Force, Ching-Ling Lien, Takako Makita, Aldons J Lusis, S Ram Kumar, Henry M Sucov

PubMed link: 28783163

Funding Grants: CIRM Stem Cell Biology Training Program

Public Summary:

Scientific Abstract:

Adult mammalian cardiomyocyte regeneration after injury is thought to be minimal. Mononuclear diploid cardiomyocytes (MNDCMs), a relatively small subpopulation in the adult heart, may account for the observed degree of regeneration, but this has not been tested. We surveyed 120 inbred mouse strains and found that the frequency of adult mononuclear cardiomyocytes was surprisingly variable (>7-fold). Cardiomyocyte proliferation and heart functional recovery after coronary artery ligation both correlated with pre-injury MNDCM content. Using genome-wide association, we identified *Tnni3k* as one gene that influences variation in this composition and demonstrated that *Tnni3k* knockout resulted in elevated MNDCM content and increased cardiomyocyte proliferation after injury. Reciprocally, overexpression of *Tnni3k* in zebrafish promoted cardiomyocyte polyploidization and compromised heart regeneration. Our results corroborate the relevance of MNDCMs in heart regeneration. Moreover, they imply that intrinsic heart regeneration is not limited nor uniform in all individuals, but rather is a variable trait influenced by multiple genes.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/frequency-mononuclear-diploid-cardiomyocytes-underlies-natural-variation>